

Reg. No. : MG122CCS.R.25.

Name : Shainima C.K.....

**I Semester B.Sc. Degree (C.B.C.S.S. – O.B.E. – Regular/Supplementary/
Improvement) Examination, November 2022**
(2019 Admission Onwards)

COMPLEMENTARY ELECTIVE COURSE IN MATHEMATICS
1C01 MAT-CS : Mathematics for Computer Science – I

Time : 3 Hours

Max. Marks : 40

PART – A

Answer any four questions from this Part. Each question carries 1 mark. (4×1=4)

1. Find $D^n(ax + b)^m$.
2. Find the Maclaurin's series expansion of the function $\cos x$.
3. Evaluate $\lim_{x \rightarrow 0} \frac{\log x}{\cot x}$.
4. Define rank of a matrix.
5. If the rank of the matrix $\begin{pmatrix} 12 & 9 \\ y & 3 \end{pmatrix}$ is one, then find y.

PART – B

Answer any 7 questions from this Part. Each question carries 2 marks. (7×2=14)

6. If $x = a(\cos t + t \sin t)$, $y = a(\sin t - t \cos t)$, find $\frac{d^2y}{dx^2}$.
7. Find $D^n[e^{ax} \cos (bx + c)]$.
8. Find the n^{th} derivative of $e^x(2x + 3)^3$.
9. Verify Cauchy's Mean-value theorem for the function e^x and e^{-x} in the interval (a, b) .
10. Verify Rolle's theorem for $f(x) = (x + 2)^2 (x - 3)^4$ in $(-2, 3)$.

K22U 3424



11. Evaluate $\lim_{x \rightarrow 0} \left(\frac{\tan x}{3} \right)^{\frac{1}{x^2}}$.

12. Are the vectors $x_1 = (1, 3, 4, 2)$, $x_2 = (3, -5, 2, 2)$, $x_3 = (2, -1, 3, 2)$ linearly dependent? If so express one of these as a linear combination of the others?

13. Using Gauss-Jordan method, find the inverse of the matrix $\begin{pmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{pmatrix}$.

14. Show that the transformation $y_1 = 2x_1 + x_2 + x_3$, $y_2 = x_1 + x_2 + 2x_3$, $y_3 = x_1 - 2x_2$ is regular. Write down the inverse transformation.

15. Reduce the law $y = mx^n + c$ into a linear law.

PART - C

Answer any 4 questions from this Part. Each question carries 3 marks. (4x3)

16. If $x^3 + y^3 = 3axy$, prove that $\frac{d^2y}{dx^2} = \frac{2a^2xy}{(y^2 - ax)^3}$.

17. Find the n^{th} derivative of $\frac{x}{(x-1)(2x+3)}$.

18. Prove that (if $0 < a < b < 1$), $\frac{b-a}{1+b^2} < \tan^{-1} b - \tan^{-1} a < \frac{b-a}{1+a^2}$. Hence show that $\frac{\pi}{4} + \frac{3}{25} < \tan^{-1} \frac{4}{3} < \frac{\pi}{4} + \frac{1}{6}$.

19. Expand $e^{a \sin^{-1} x}$ in ascending powers of x .

20. Using partition method, find the inverse of $\begin{pmatrix} 1 & 1 & 1 \\ 4 & 3 & -1 \\ 3 & 5 & 3 \end{pmatrix}$.

21. Test for consistency and solve

$$5x + 3y + 7z = 4, 3x + 26y + 2z = 9, 7x + 10y + 2z = 5.$$

22. Write the working procedure to fit the straight line $y = a + bx$ to a given data.

PART - D

Answer **any 2** questions from this Part. **Each** question carries 5 marks. $(2 \times 5 = 10)$

23. If $y = e^{a\sin^{-1}x}$, prove that $(1 - x^2)y_{n+2} - (2n + 1)xy_{n+1} - (n^2 + a^2)y_n = 0$. Hence find that value of y_n when $x = 0$.

24. Evaluate $\lim_{x \rightarrow 0} \frac{(1+x)^{\frac{1}{x}} - e}{x}$.

25. Find that value of λ for which the equations

$\checkmark (\lambda - 1)x + (3\lambda + 1)y + 2\lambda z = 0$

$(\lambda - 1)x + (4\lambda - 2)y + (\lambda + 3)z = 0$

$2x + (3\lambda + 1)y + 3(\lambda - 1)z = 0$.

are consistent and find the ratios of $x : y : z$ when λ has the smallest of these values. What happens when λ has the greatest of these values.

26. Fit a second degree parabola to the following data :

\checkmark

x	1989	1990	1991	1992	1993	1994	1995	1996	1997
---	------	------	------	------	------	------	------	------	------

\checkmark

y	352	356	357	358	360	361	361	360	359
---	-----	-----	-----	-----	-----	-----	-----	-----	-----